

# Erdosteine

## Zertin<sup>®</sup> Oral Preparations

### FORMULATION

*Erdosteine (Zertin<sup>®</sup>) 300mg Capsule:* With yellow body and green cap, each capsule contains 300mg Erdosteine.

*Erdosteine (Zertin<sup>®</sup>) 175mg/5mL Powder for Suspension:* Free flowing white granulate with characteristic pleasant odor and orange flavour. Each 5mL of reconstituted suspension contains 175mg Erdosteine.

### INDICATIONS

Treatment of patients with acute and chronic bronchopulmonary diseases, rhinosinusitis, laryngopharyngitis or exacerbations of these chronic diseases in association with mucus production and mucus transport.

### DOSAGE & ADMINISTRATION

#### Adults

300mg Capsule : One capsule twice daily.  
175mg/5mL Powder for Suspension : 8.5mL (300mg) twice daily.

No dose adjustment is required in treatment of adult relapses of COPD, or in geriatric patients.

#### Paediatric use:

As a general rule, 10mg/Kg/day in two administrations at 12 hours intervals.

175mg/5mL Powder for Suspension : the recommended dosage regimen is:

Body Weight (Kg)	Age	Dose
10 – 20 Kg	2 – 6 years	2.5 mL twice daily
21 – 30 Kg	7 – 12 years	5.0 mL twice daily
>30 Kg	> 12 years	5.0 mL three times daily (or 7.5mL twice daily)

#### Directions for Reconstitution:

To make 60mL reconstituted suspension, loosen the powder by shaking and slowly add 42mL of water, then replace cap and shake well.

To make 100mL reconstituted suspension, loosen the powder by shaking and slowly add 70mL of water, then replace cap and shake well.

Always shake well before use. The reconstituted suspension can be stored in a refrigerator for a maximum of 10 days.

### CONTRAINDICATIONS

Hypersensitivity to the active substance or to any of the excipients or to free SH-group containing products.

Hepatic disorders and abnormalities (e.g. increase of serum alkaline phosphatase, transaminases, etc.).

Renal insufficiency (creatinine clearance <25mL/min).

Homocystinuria (the active substance is partially metabolised to homocysteine and there are no data concerning administration of erdosteine in case of congenital errors of the metabolism of aminoacids, especially in those patients obliged to follow a methionine-free dietary regimen).

Phenylketonuria (only for powder for suspension due to the presence of aspartame in this presentation).

### WARNINGS & PRECAUTIONS

In case of appearance of classical hypersensitivity signs and symptoms, the treatment with erdosteine must be immediately suspended.

The simultaneous administration of an anti-cough treatment is irrational and could induce the accumulation of liquefied excreatum in the bronchial tree with an increase of risk of superinfection or bronchospasm.

Safety and efficacy in children under two years of age have not been established.

In case of co-treatment with other drugs see the individual product data sheet.

### **Drug Interaction**

No harmful interaction has been observed with other drugs, often used in treatment of respiratory infections and COPD, like theophylline, bronchodilating compounds (corticoid treatment), erythromycin, amoxicillin or co-trimoxazole.

Erdosteine increases amoxicillin concentration in the respiratory tract.

The simultaneous administration of an anti-cough treatment is irrational and could induce the accumulation of liquefied excreatum in the bronchial tree with an increase of risk of superinfection or bronchospasm.

### **Pregnancy and Lactation**

No embryo-foetal abnormalities have been observed in pre-clinical studies.

Information of erdosteine use during pregnancy and nursing is limited. Therefore erdosteine should be administered to pregnant or nursing women only when considered necessary by the clinician.

### **Effects on the ability to drive and use machines**

No effects on the ability to drive and use machines have been observed.

### **Incompatibilities**

None known.

### **ADVERSE EVENTS**

Administration of erdosteine could occasionally cause gastro-intestinal side-effect, like gastric burning, nausea and rarely diarrhea.

In few cases, at the beginning of treatment, ageusia or dysgeusia have been observed.

Hypersensitivity reactions (skin rash, urticaria, unexpected hyperpyrexia) are rare.

The following adverse reaction rates are based on the use of erdosteine in treated population (1520 patients) during phase III clinical trials:

1 to 3%	:	gastralgia, nausea, headache
0.5 to 1%	:	constipation, diarrhea, mouth dryness, vertigo, general malaise.

The observed adverse reactions are not different, from a qualitative point of view, from those observed under placebo.

### **OVERDOSAGE**

No case of overdosage has been reported.

### **CLINICAL PHARMACOLOGY**

#### **Pharmacodynamics**

*Erdosteine (Zertin<sup>®</sup>)* Capsule and Powder for Suspension contain erdosteine, an original derivative of natural mercapto-aminoacid in thiolactonic form. Following oral administration erdosteine is rapidly metabolised in the liver. The product acts as a pro-drug and its metabolites are mainly responsible for mucolytic activity, due to the presence of free thiol groups which cause the splitting up of the intra- and inter-molecular disulphide bridges of several proteins and mucoproteins present in the expectoration, resulting in a reduction of the mucus elasticity and viscosity.

As a consequence, the reduction of mucus viscosity allows an improved penetration of amoxicillin in the mucus itself, without effects on the amoxicillin specific activity or on the plasmatic levels of amoxicillin and erdosteine. At the same time, the mucociliary transport of mucus improves.

The free thiol groups of erdosteine metabolites inactive oxidizing substances, in particular oxygen radicals, giving erdosteine ability to prevent:

- oxidation of  $\alpha_1$ -antitrypsin (which in its reduced form possesses elastase inhibitory activity)
- reduction of the chemotactic activity of polynuclear neutrophils, caused by cigarette smoke
- smoke induced antipyrine oxidation.

Following erdosteine administration, increased plasmatic and bronchoalveolar lavage (BAL) levels of reduced glutathione (GSH) are observed.

Erdosteine has been shown to increase respiratory tract IgA concentration in COPD patients and to prevent the inhibition of granulocytes chemotaxis caused by smoking.

Erdosteine increases the concentration of amoxicillin in the bronchial secretions and co-administration gives a faster response in comparison with mono-therapy with amoxicillin.

The presence of free SH-group in erdosteine metabolites results in reduced bacterial adhesiveness to human mucosal cells, leading to a reduced bacterial virulence.

As in erdosteine, per se, the thiol groups are masked, there are little effects on the GI tract, at recommended dose.

### **Pharmacokinetics**

Erdosteine is rapidly absorbed and metabolised at hepatic level into three main metabolites, containing free SH- groups. The most abundant and active is N-thiodiglycolyl-homocysteine (Metabolite 1, also M1).

After single dose or repeated dose the main pharmacokinetic parameters of erdosteine and its metabolites are not statistically different.

Peak plasma concentrations of Erdosteine (*Zertin*<sup>®</sup>) and metabolite 1 are achieved in approximately 1 and 3 hours, respectively.

The protein binding rate of erdosteine is about 64.5%.

Elimination occurs mainly by renal route in form of sulphates, only little amount of unmetabolised erdosteine goes through faecal excretion.

Repeated administration and food intake do not affect significantly the pharmacokinetic parameters, as  $C_{max}$  and AUC are not affected; after a meal, only  $T_{max}$  is slightly shifted.

In particular neither accumulation nor enzymatic induction has been observed.

In case of hepatic impairment an increase of  $C_{max}$  and AUC values has been observed. Moreover, in case of severe hepatic dyscrasia an increase of  $T_{1/2\beta}$  can be noticed.

In severe renal impairment (creatinine clearance 25 to 40mL/min) there is the risk of metabolites accumulation.

Similar findings result for the 175mg/5mL Powder for Suspension.

### **PRE-CLINICAL SAFETY DATA**

#### **Toxicity**

The acute toxicity of erdosteine is low, the  $LD_{50}$  values ranging from >5000mg/Kg (p.o.: mouse, rat and i.p. in rat) to >3500mg/Kg (i.v., mouse).

In sub-acute toxicity studies, erdosteine did not cause pathological alterations within the dose range of 100 – 1000mg/Kg/day in rats and of 100mg/Kg/day in dogs (p.o. for 4 weeks), up to 4500mg/Kg/day in rats exposed to aerosol (2 hrs/day) for 4 weeks. Only the highest dose in dogs (400mg/Kg/day) induced a slightly abnormal increase of liver weight with modest histological alterations.

Similarly, during long-term studies (26 weeks) no important toxic symptoms resulted following treatment p.o. with up to 1000mg/Kg/day (rats) and 200mg/Kg/day (dogs, higher dose).

Higher doses in rats induced a reversible decrease of body weight gain and blood protein levels.

No harmful effects on lungs, liver, heart or kidneys were observed.

CNS effects, in terms of sedation, hypothermia and prostration, were observed at extremely high doses (5000mg/Kg p.o., 3500mg/Kg i.v. and i.p. in rats).

The local tolerability was good following oral, inhalation and rectal administration.

#### **Reproductive function**

The higher dose (1000mg/Kg/day p.o.) does not appear to induce toxic effects on fertility and general reproductive performance in rats.

#### **Embryo-foetal and perinatal toxicity**

Erdosteine up to the daily doses of 1000mg/Kg (rats) and 700mg/Kg (rabbits) by oral route appears as lacking of foetotoxic, embryotoxic and teratogenic effects; the higher dose for rats is also devoid of any effect on the peri- and post-natal parameters.

#### **Mutagenic potential**

No mutagenic potential for Erdosteine resulted from studies according to several experimental models: gene mutation on bacteria (Ames test, gene conversion on yeast) and eucaryotes (punctiform mutation); chromosomal aberration in mammals (*in vitro*: Chinese hamster and human lymphocytes; *in vivo* micronucleus test, host mediated assay); host mediated assay and urinary assay were performed as well.

#### **Carcinogenic potential**

In view of the molecular structure of the product (a derivative of a natural aminoacid), which does not have any similarity with known carcinogenic compounds, no study was performed.

#### **AVAILABILITY**

\* *Erdosteine (Zertin®) 300mg Capsule*: Cartons of 20 capsules in blister packs of 10.

\*\* *Erdosteine (Zertin®) 175mg/5mL Powder for Suspension*: Bottles of 60mL and 100mL powder for suspension

#### **STORAGE**

*Capsule*: Store at temperatures not exceeding 25°C. Keep in a dry place.

*Powder for Suspension*: Store at temperatures not exceeding 30°C. Shake well before use.

The reconstituted suspension can be stored in a refrigerator for a maximum of 10 days.

#### **CAUTION**

Food, Drugs, Devices and Cosmetics Act prohibits dispensing without prescription.

Keep all medicines out of reach of children.

ZERTIN is a registered trademark of the GlaxoSmithKline group of companies.

All rights reserved.

© 2011 GlaxoSmithKline group of companies.

Version number: I-ZYX-2

Revision date: 17 October 2011



GlaxoSmithKline Philippines Inc  
2266 Chino Roces Avenue, City of Makati

\* Mfd by: Famar Italia, S.p.A., Milan, Italy for  
Edmond Pharma S.R.L., Milan, Italy

\*\*Mfd. by: Hizon Laboratories, Inc.  
Assumption Road, Sumulong Highway, Antipolo City  
Under license from: Edmond Pharma S.R.L., Milan, Italy

PH/ZER/0020/15

August 2015

For Healthcare Professionals Only

ZERTIN is a registered trademark of GSK group of companies